


## Arc's Midpoint Turns Kinetic, Adds Applied to Theoretic...

Gregory V. Akulov, teacher, Luther College High School, Regina, Saskatchewan

[auf Duetsch](#)

[en français](#)

An object, initially at rest, slides from the top of a vertical semicircular path with a horizontal diameter (see diagram). At points **A**, **B**, **C**, and **M** along the semicircle, the object has kinetic energy in joules  $a = 62 \text{ J}$ ,  $b = 638 \text{ J}$ ,  $c = 962 \text{ J}$ , and  $\mu \text{ J}$ , respectively. If **C** is the lowest point of the path, and **M** is equidistant from **A** and **B**, determine the **exact value** of  $\mu$ . Assume that the path is frictionless and the air resistance is negligible.



The [arc midpoint computation](#)<sup>1</sup> approach to solving this problem finds

$$2\mu = \sqrt{(962 \text{ J} + 638 \text{ J})(962 \text{ J} + 62 \text{ J})} + \sqrt{(962 \text{ J} - 638 \text{ J})(962 \text{ J} - 62 \text{ J})}, \text{ and } \mu = 910 \text{ J}$$

**1. Verify** the answer using alternative approach. **Compare** solutions.

**2. Show** that the values of **kinetic energy**  $a$ ,  $b$ ,  $c$ , and  $\mu$  satisfy

$$2\mu = \sqrt{(c + a)(c + b)} \pm \sqrt{(c - a)(c - b)} \quad (1)$$

**3. Specify** when (1) requires the **sum** of radicals, and when it requires their **difference**.

Copyright © August 8, 2018 by Gregory V. Akulov

<sup>1</sup> <http://mathcentral.uregina.ca/RR/database/RR.09.10/akulov2.html>